


Graphs and Linear Functions

A 2-dimensional graph is a visual representation of a relationship between two variables given by an equation or an inequality. Graphs help us solve algebraic problems by analysing the geometric aspects of a problem. While equations are more suitable for precise calculations, graphs are more suitable for showing patterns and trends in a relationship. To fully utilize what graphs can offer, we must first understand the concepts and skills involved in graphing that are discussed in this chapter.



G1

System of Coordinates, Graphs of Linear Equations and the Midpoint Formula

In this section, we will review the rectangular coordinate system, graph various linear equations and inequalities, and introduce a formula for finding coordinates of the midpoint of a given segment.

The Cartesian Coordinate System




Figure 1a




Figure 1b

A rectangular coordinate system, also called a **Cartesian coordinate system** (in honor of French mathematician, **René Descartes**), consists of two perpendicular number lines that cross each other at point zero, called the **origin**. Traditionally, one of these number lines, usually called the **x-axis**, is positioned horizontally and directed to the right (see *Figure 1a*). The other number line, usually called **y-axis**, is positioned vertically and directed up. Using this setting, we identify each point P of the plane with an **ordered pair** of numbers (x, y) , which indicates the location of this point with respect to the origin. The first number in the ordered pair, the **x-coordinate**, represents the horizontal distance of the point P from the origin. The second number, the **y-coordinate**, represents the vertical distance of the point P from the origin. For example, to locate point $P(3,2)$, we start from the origin, go 3 steps to the right, and then two steps up. To locate point $Q(-3, -2)$, we start from the origin, go 3 steps to the left, and then two steps down (see *Figure 1b*).

Observe that the coordinates of the origin are $(0,0)$. Also, the second coordinate of any point on the x -axis as well as the first coordinate of any point on the y -axis is equal to zero. So, points on the x -axis have the form $(x, 0)$, while points on the y -axis have the form of $(0, y)$.

To **plot** (or **graph**) an ordered pair (x, y) means to place a dot at the location given by the ordered pair.

Example 1



Plotting Points in a Cartesian Coordinate System

Plot the following points:


$$A(2, -3), \quad B(0, 2), \quad C(1, 4), \quad D(-5, 0),$$

$$E(-2, -3), \quad F(0, -4), \quad G(-3, 3)$$

Solution



Remember! The order of numbers in an ordered pair is important! The **first** number represents the **horizontal** displacement and the **second** number represents the **vertical** displacement from the origin.



Graphs of Linear Equations

A **graph of an equation** in two variables, x and y , is the set of points corresponding to **all ordered pairs (x, y)** that **satisfy** the equation (make the equation true). This means that a graph of an equation is the visual representation of the **solution set** of this equation.

To determine if a point (a, b) belongs to the graph of a given equation, we check if the equation is satisfied by $x = a$ and $y = b$.

Example 2 ▶ Determining if a Point is a Solution of a Given Equation

Determine if the points $(5, 3)$ and $(-3, -2)$ are solutions of $2x - 3y = 0$.

Solution ▶ After substituting $x = 5$ and $y = 3$ into the equation $2x - 3y = 0$, we obtain

$$\begin{aligned} 2 \cdot 5 - 3 \cdot 3 &= 0 \\ 10 - 9 &= 0 \\ 1 &= 0, \end{aligned}$$

which is not true. Since the coordinates of the point $(5, 3)$ do not satisfy the given equation, the point **$(5, 3)$ is not a solution** of this equation.

Note: The fact that the point $(5, 3)$ **does not satisfy** the given equation indicates that it **does not belong to the graph** of this equation.

However, after substituting $x = -3$ and $y = -2$ into the equation $2x - 3y = 0$, we obtain

$$\begin{aligned} 2 \cdot (-3) - 3 \cdot (-2) &= 0 \\ -6 + 6 &= 0 \\ 0 &= 0, \end{aligned}$$

which is true. Since the coordinates of the point $(-3, -2)$ satisfy the given equation, the point **$(-3, -2)$ is a solution** of this equation.

Note: The fact that the point $(-3, -2)$ **satisfies** the given equation indicates that it **belongs to the graph** of this equation.

To find a solution to a given equation in two variables, we choose a particular value for one of the variables, substitute it into the equation, and then solve the resulting equation for the other variable.

For example, to find a solution to $3x + 2y = 6$, we can choose for example $x = 0$, which leads us to

$$\begin{aligned} 3 \cdot 0 + 2y &= 6 \\ 2y &= 6 \\ y &= 3. \end{aligned}$$

This means that the point **$(0, 3)$** satisfies the equation and therefore belongs to the graph of this equation. If we choose a different x -value, for example $x = 1$, the corresponding y -value becomes

$$\begin{aligned} 3 \cdot 1 + 2y &= 6 \\ 2y &= 3 \\ y &= \frac{3}{2} \end{aligned}$$

So, the point $\left(1, \frac{3}{2}\right)$ also belongs to the graph.

Since any real number could be selected for the x -value, there are infinitely many solutions to this equation. Obviously, we will not be finding all of these infinitely many ordered pairs of numbers in order to graph the solution set to an equation. Rather, based on the location of several solutions that are easy to find, we will look for a pattern and predict the location of the rest of the solutions to complete the graph.

To find more points that belong to the graph of the equation in our example, we might want to solve $3x + 2y = 6$ for y . The equation is equivalent to

$$\begin{aligned} 2y &= -3x + 6 \\ y &= -\frac{3}{2}x + 3 \end{aligned}$$

Observe that if we choose x -values to be multiples of 2, the calculations of y -values will be easier in this case. Here is a table of a few more (x, y) points that belong to the graph:

x	$y = -\frac{3}{2}x + 3$	(x, y)
-2	$-\frac{3}{2}(-2) + 3 = 6$	$(-2, 6)$
2	$-\frac{3}{2}(2) + 3 = 0$	$(2, 0)$
4	$-\frac{3}{2}(4) + 3 = -3$	$(4, -3)$




Figure 2a




Figure 2b

After plotting the obtained solutions, $(-2, 6)$, $(0, 3)$, $\left(1, \frac{3}{2}\right)$, $(2, 0)$, $(4, -3)$, we observe that the points appear to lie on the same line (see *Figure 2a*). If all the ordered pairs that satisfy the equation $3x + 2y = 6$ were graphed, they would form the line shown in *Figure 2b*. Therefore, if we knew that the graph would turn out to be a line, it would be enough to find just two points (solutions) and draw a line passing through them.

How do we know whether or not the graph of a given equation is a line? It turns out that:

For any equation in two variables, the graph of the equation is a **line** if and only if (iff) the equation is **linear**.

So, the question is how to recognize a linear equation?

Definition 1.1 ▶ Any equation that can be written in the form

$$Ax + By = C, \text{ where } A, B, C \in \mathbb{R}, \text{ and } A \text{ and } B \text{ are not both } 0,$$

is called a **linear equation** in two variables.

The form $Ax + By = C$ is called **standard form** of a linear equation.

Example 3 ▶ **Graphing Linear Equations Using a Table of Values**


Graph $4x - 3y = 6$ using a table of values.

Solution ▶ Since this is a linear equation, we expect the graph to be a line. While finding two points satisfying the equation is sufficient to graph a line, it is a good idea to use a third point to guard against errors. To find several solutions, first, let us solve $4x - 3y = 6$ for y :

$$\begin{aligned} -3y &= -4x + 6 \\ y &= \frac{4}{3}x - 2 \end{aligned}$$

We like to choose x -values that will make the calculations of the corresponding y -values relatively easy. For example, if x is a multiple of 3, such as -3 , 0 or 3 , the denominator of $\frac{4}{3}$ will be reduced. Here is the table of points satisfying the given equation and the graph of the line.

x	$y = \frac{4}{3}x - 2$	(x, y)
-3	$\frac{4}{3}(-3) - 2 = -6$	$(-3, -6)$
0	$\frac{4}{3}(0) - 2 = -2$	$(0, -2)$
3	$\frac{4}{3}(3) - 2 = 2$	$(3, 2)$



To graph a linear equation in standard form, we can develop a table of values as in *Example 2*, or we can use the x - and y -intercepts.

Definition 1.2 ▶ The **x -intercept** is the point (if any) where the graph intersects the x -axis. So, the x -intercept has the form $(x, 0)$.

The **y -intercept** is the point (if any) where the graph intersects the y -axis. So, the y -intercept has the form $(0, y)$.

Example 4 ▶ **Graphing Linear Equations Using x - and y -intercepts**

Graph $5x - 3y = 15$ by finding and plotting the x - and y -intercepts.

Solution ▶ To find the x -intercept, we substitute $y = 0$ into $5x - 3y = 15$, and then solve the resulting equation for y . So, we have

$$\begin{aligned}5x &= 15 \\x &= 3.\end{aligned}$$

To find y -intercept, we substitute $x = 0$ into $5x - 3y = 15$, and then solve the resulting equation for x . So,


$$\begin{aligned}-3y &= 15 \\y &= -5.\end{aligned}$$

Hence, we have

x -intercept

y -intercept

x	y
3	0
0	-5



To find several points that belong to the graph of a linear equation in two variables, it was easier to solve the standard form $Ax + By = C$ for y , as follows

$$\begin{aligned}By &= -Ax + C \\y &= -\frac{A}{B}x + \frac{C}{B}.\end{aligned}$$

This form of a linear equation is also very friendly for graphing, as the graph can be obtained without any calculations. See *Example 4*.

Any equation $Ax + By = C$, where $B \neq 0$ can be written in the form

$$y = mx + b,$$

which is referred to as the **slope-intercept form** of a linear equation.

The value $m = -\frac{A}{B}$ represents the **slope** of the line. Recall that **slope** = $\frac{\text{rise}}{\text{run}}$.

The value b represents the y -intercept, so the point $(0, b)$ belongs to the graph of this line.

Example 5 ▶ Graphing Linear Equations Using Slope and y -intercept

Determine the slope and y -intercept of each line and then graph it.

a. $y = \frac{2}{3}x + 1$


b. $5x + 2y = 8$

Solution ▶

a. The slope is the coefficient by x , so it is $\frac{2}{3}$.

The y -intercept is $(0, 1)$.

So we plot point $(0, 1)$ and then, since $\frac{2}{3} = \frac{\text{rise}}{\text{run}}$, we rise 2 units and run 3 units to find the next point that belongs to the graph.




- b. To see the slope and y -intercept, we solve $5x + 2y = 8$ for y .

$$2y = -5x + 8$$

$$y = \frac{-5}{2}x + 4$$


So, the slope is $\frac{-5}{2}$ and the y -intercept is $(0,4)$. We start from $(0,4)$ and then run 2 units and fall 5 units (because of -5 in the numerator).



Note: Although we can *run* to the right or to the left, depending on the sign in the denominator, we usually **keep the denominator positive and always run forward** (to the right). If the slope is negative, we **keep the negative sign in the numerator** and either *rise* or *fall*, depending on this sign. However, when finding additional points of the line, sometimes we can repeat the *run/rise* movement in either way, to the right, or to the left from one of the already known points. For example, in *Example 4a*, we could find the additional point at $(-3, -1)$ by *running* 3 units to the left and 2 units down from $(0, 1)$, as the slope $\frac{2}{3}$ can also be seen as $\frac{-2}{-3}$, if needed.


Some linear equations contain just one variable. For example, $x = 3$ or $y = -2$. How would we graph such equations in the xy -plane?

Observe that $y = -2$ can be seen as $y = 0x - 2$, so we can graph it as before, using the **slope** of **zero** and the **y -intercept** of $(0, -2)$. The graph consists of all points that have y -coordinates equal to -2 . Those are the points of type $(x, -2)$, where x is any real number. The graph is a **horizontal line** passing through the point $(0, -2)$.



Note: The horizontal line $y = 0$ is the x -axis.

The equation $x = 3$ doesn't have a slope-intercept representation, but it is satisfied by any point with x -coordinate equal to 3. So, by plotting several points of the type $(3, y)$, where y is any real number, we obtain a **vertical line** passing through the point $(3, 0)$. This particular line doesn't have a y -intercept, and its **slope** = $\frac{\text{rise}}{\text{run}}$ is considered to be **undefined**. This is because the "*run*" part calculated between any two points on the line is equal to zero and we can't perform division by zero.



Note: The vertical line $x = 0$ is the y -axis.

In general, the graph of any equation of the type

$$y = b, \text{ where } b \in \mathbb{R}$$

is a **horizontal line** with y -intercept $(0, b)$. The **slope** of such a line is **zero**.

The graph of any equation of the type

$$x = a, \text{ where } a \in \mathbb{R}$$

is a **vertical line** with x -intercept $(a, 0)$. The **slope** of such a line is **undefined**.

Example 6 ▶ Graphing Special Types of Linear Equations

Graph each equation and state its slope.



a. $x = -1$

b. $y = 0$

c. $y = x$


Solution ▶

- a. The solutions to the equation $x = -1$ are all pairs of the type $(-1, y)$, so after plotting points like $(-1, 0)$, $(-1, 2)$, etc., we observe that the graph is a **vertical line** intercepting the x -axis at $x = -1$. So the **slope** of this line is **undefined**.



- b. The solutions to the equation $y = 0$ are all pairs of the type $(x, 0)$, so after plotting points like $(0, 0)$, $(0, 3)$, etc., we observe that the graph is a **horizontal line** following the x -axis. The **slope** of this line is **zero**.

- c. The solutions to the equation $y = x$ are all pairs of the type (x, x) , so after plotting points like $(0, 0)$, $(2, 2)$, etc., we observe that the graph is a **diagonal line**, passing through the origin and making a 45° angle with the x -axis. The **slope** of this line is **1**.





Observation: A graph of any equation of the type $y = mx$ is a line passing through the origin, as the point $(0, 0)$ is one of the solutions.

Midpoint Formula

To find a representative value of a list of numbers, we often calculate the average of these numbers. Particularly, to find an average of, for example, two test scores, 72 and 84, we take half of the sum of these scores. So, the average of 72 and 84 is equal to $\frac{72+84}{2} = \frac{156}{2} = 78$. Observe that 78 lies on a number line exactly halfway between 72 and 84. The idea of taking an average is employed in calculating coordinates of the midpoint of any line segment.

Definition 1.3 ▶ The **midpoint** of a line segment is the point of the segment that is equidistant from both ends of this segment.



Suppose $A = (x_1, y_1)$, $B = (x_2, y_2)$, and M is the **midpoint** of the line segment \overline{AB} . Then the x -coordinate of M lies halfway between the two end x -values, x_1 and x_2 , and the y -coordinate of M lies halfway between the two end y -values, y_1 and y_2 . So, the coordinates of the midpoint are **averages** of corresponding x -, and y -coordinates:

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) \quad (1)$$

Example 7 ▶ **Finding Coordinates of the Midpoint**

Find the midpoint M of the line segment connecting $P = (-3, 7)$ and $Q = (5, -12)$.


Solution ▶ The coordinates of the midpoint M are averages of the x - and y -coordinates of the endpoints. So,

$$M = \left(\frac{-3 + 5}{2}, \frac{7 + (-12)}{2} \right) = \left(1, -\frac{5}{2} \right).$$

Example 8 ▶ **Finding Coordinates of an Endpoint Given the Midpoint and the Other Endpoint**

Suppose segment AB has its midpoint M at $(2, 3)$. Find the coordinates of point A , knowing that $B = (5, 6)$.

Solution ▶ Let $A = (x, y)$ and $B = (5, 6)$. Since $M = (2, 3)$ is the midpoint of \overline{AB} , by formula (1), the following equations must hold:



$$\frac{x+5}{2} = 2 \quad \text{and} \quad \frac{y+6}{2} = 3$$

Multiplying these equations by 2, we obtain

$$x + 5 = 4 \quad \text{and} \quad y + 6 = 6,$$


which results in

$$x = -1 \quad \text{and} \quad y = 0.$$

Hence, the coordinates of point A are $(-1, 0)$.

G.1 Exercises

- Plot each point in a rectangular coordinate system.
 - $(1, 2)$
 - $(-2, 0)$
 - $(0, -3)$
 - $(4, -1)$
 - $(-1, -3)$
- State the coordinates of each plotted point.



Determine if the given ordered pair is a solution of the given equation.

- $(-2, 2)$; $y = \frac{1}{2}x + 3$
- $(4, -5)$; $3x + 2y = 2$
- $(5, 4)$; $4x - 5y = 1$

Graph each equation using the suggested table of values.

- $y = 2x - 3$
- $y = -\frac{1}{3}x + 2$
- $x + y = 3$
- $4x - 5y = 20$

x	y
0	
1	
2	
3	

x	y
-3	
0	
3	
6	

x	y
0	
	0
-1	
	1

x	y
0	
	0
2	
	-3

Graph each equation using a table of values.

- $y = \frac{1}{3}x$
- $y = \frac{1}{2}x + 2$
- $6x - 3y = -9$
- $6x + 2y = 8$
- $y = \frac{2}{3}x - 1$
- $y = -\frac{3}{2}x$
- $3x + y = -1$
- $2x = -5y$
- $-3x = -3$
- $6y - 18 = 0$
- $y = -x$
- $2y - 3x = 12$

Determine the x - and y -intercepts of each line and then graph it. Find additional points, if needed.

- $5x + 2y = 10$
- $x - 3y = 6$
- $8y + 2x = -4$
- $3y - 5x = 15$
- $y = -\frac{2}{5}x - 2$
- $y = \frac{1}{2}x - \frac{3}{2}$
- $2x - 3y = -9$
- $2x = -y$

Determine the **slope** and **y-intercept** of each line and then graph it.

30. $y = 2x - 3$

31. $y = -3x + 2$

32. $y = -\frac{4}{3}x + 1$

33. $y = \frac{2}{5}x + 3$

34. $2x + y = 6$

35. $3x + 2y = 4$

36. $-\frac{2}{3}x - y = 2$

37. $2x - 3y = 12$

38. $2x = 3y$

39. $y = \frac{3}{2}$

40. $y = x$

41. $x = 3$

Find the midpoint of each segment with the given endpoints.

42. $(-8, 4)$ and $(-2, -6)$

43. $(4, -3)$ and $(-1, 3)$

44. $(-5, -3)$ and $(7, 5)$

45. $(-7, 5)$ and $(-2, 11)$

46. $(\frac{1}{2}, \frac{1}{3})$ and $(\frac{3}{2}, -\frac{5}{3})$

47. $(\frac{3}{5}, -\frac{1}{3})$ and $(\frac{1}{2}, -\frac{5}{2})$

Segment AB has the given coordinates for the endpoint A and for its midpoint M . Find the coordinates of the endpoint B .

48. $A(-3, 2), M(3, -2)$

49. $A(7, 10), M(5, 3)$

50. $A(5, -4), M(0, 6)$

51. $A(-5, -2), M(-1, 4)$

G2

Slope of a Line and Its Interpretation

Slope (steepness) is a very important concept that appears in many branches of mathematics as well as statistics, physics, business, and other areas. In algebra, slope is used when graphing lines or analysing linear equations or functions. In calculus, the concept of slope is used to describe the behaviour of many functions. In statistics, slope of a regression line explains the general trend in the analysed set of data. In business, slope plays an important role in linear programming. In addition, slope is often used in many practical ways, such as the slope of a road (*grade*), slope of a roof (*pitch*), slope of a ramp, etc. In this section, we will define, calculate, and provide some interpretations of slope.



Slope




Figure 1a

Given two lines, a and b , how can we tell which one is steeper? One way to compare the steepness of these lines is to move them closer to each other, so that a point of intersection, P , can be seen, as in *Figure 1a*. Then, after running horizontally a few steps from the point P , draw a vertical line to observe how high the two lines have risen. The line that crosses this vertical line at a higher point is steeper. So, for example in *Figure 1a*, line a is steeper than line b . Observe that because we run the same horizontal distance for both lines, we could compare the steepness of the two lines just by looking at the vertical rise. However, since the *run* distance can be chosen arbitrarily, to represent the steepness of any line, we must look at the *rise* (vertical change) in respect to the *run* (horizontal change). This is where the concept of slope as a ratio of *rise* to *run* arises.




Figure 1b

To measure the slope of a line or a line segment, we choose any two distinct points of such a figure and calculate the ratio of the **vertical change** (*rise*) to the **horizontal change** (*run*) between the two points. For example, the slope between points $A(1,2)$ and $B(3,5)$ equals

$$\frac{\text{rise}}{\text{run}} = \frac{3}{2},$$

as in *Figure 1a*. If we rewrite this ratio so that the denominator is kept as one,

$$\frac{3}{2} = \frac{1.5}{1} = 1.5,$$

we can think of slope as of the **rate of change in y -values with respect to x -values**. So, a slope of 1.5 tells us that the y -value increases by 1.5 units per every increase of one unit in x -value.




Figure 1c

Generally, the slope of a line passing through two distinct points, (x_1, y_1) and (x_2, y_2) , is the **ratio** of the change in y -values, $y_2 - y_1$, to the change in x -values, $x_2 - x_1$, as presented in *Figure 1c*. Therefore, the formula for calculating slope can be presented as

$$\frac{\text{rise}}{\text{run}} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\Delta y}{\Delta x},$$

where the Greek letter Δ (delta) is used to denote the change in a variable.

Definition 2.1 ▶ Suppose a line passes through two distinct points (x_1, y_1) and (x_2, y_2) .




If $x_1 \neq x_2$, then the **slope** of this line, often denoted by m , is equal to

$$m = \frac{\text{rise}}{\text{run}} = \frac{\text{change in } y}{\text{change in } x} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\Delta y}{\Delta x}$$

If $x_1 = x_2$, then the **slope** of the line is said to be **undefined**.


Example 1 ▶ **Determining Slope of a Line, Given Its Graph**

Determine the slope of each line.




Solution ▶

- a. To read the slope we choose two distinct points with integral coefficients (often called **lattice points**), such as the points suggested in the graph. Then, starting from the first point $(-2, 1)$ we **run** 5 units and **rise** 3 units to reach the second point $(3, 4)$. So, the slope of this line is $m = \frac{3}{5}$.




- b. This is a horizontal line, so the **rise** between any two points of this line is zero. Therefore the slope of such a line is also **zero**.

- c. If we refer to the lattice points $(-3, 0)$ and $(0, -1)$, then the **run** is 3 and the **rise** (or rather **fall**) is -1 . Therefore the slope of this line is $m = -\frac{1}{3}$.



run = 0 so
 $m = \text{undefined}$



Observation:

A line that **increases** from left to right has a **positive slope**.

A line that **decreases** from left to right has a **negative slope**.

The slope of a **horizontal** line is **zero**.


The slope of a **vertical** line is **undefined**.



Example 2 ▶ **Graphing Lines, Given Slope and a Point**

Graph the line with slope $-\frac{3}{2}$ that passes through the point $(-3, 4)$.

Solution ▶ First, plot the point $(-3, 4)$. To find another point that belongs to this line, start at the plotted point and run 2 units, then fall 3 units. This leads us to point $(-1, 1)$. For better precision, repeat the movement (two across and 3 down) to plot one more point, $(1, -2)$. Finally, draw a line connecting the plotted points.

**Example 3** ▶ **Calculating Slope of a Line, Given Two Points**

Determine the slope of a line passing through the points $(-3, 5)$ and $(7, -11)$.

Solution ▶ The slope of the line passing through $(-3, 5)$ and $(7, -11)$ is the quotient

$$\frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - (-11)}{-3 - 7} = \frac{5 + 11}{-10} = -\frac{16}{10} = -1.6$$

Example 4 ▶ **Determining Slope of a Line, Given Its Equation**

Determine the slope of a line given by the equation $2x - 5y = 7$.

Solution ▶ To see the slope of a line in its equation, we change the equation to its slope-intercept form, $y = mx + b$. The slope is the coefficient m . When solving $2x - 5y = 7$ for y , we obtain

$$-5y = -2x + 7$$

$$y = \frac{2}{5}x - \frac{7}{5}$$

So, the slope of this line is equal to $\frac{2}{5}$.

Example 5 ▶ **Interpreting Slope as an Average Rate of Change**

The value of a particular stock has increased from \$156.60 on February 10, 2018, to \$187.48 on November 10, 2018. What is the average rate of change of the value of this stock per month for the given period of time?



Solution ▶ The average value of the stock has increased by $187.48 - 156.60 = 30.88$ dollars over the 9 months (from February 10 to November 10). So, the slope of the line segment connecting the the values of the stock on these two days (as marked on the above chart) equals

$$\frac{30.88}{9} \cong 3.43 \text{ \$/month}$$

This means that the value of the stock was increasing on average by 3.43 dollars per month between February 10, 2018, and November 10, 2018.

Observe that the change in value was actually different in each month. Sometimes the change was larger than the calculated slope, but sometimes the change was smaller or even negative. However, the **slope** of the above segment gave us the information about the **average rate of change** in the stock's value during the stated period.

Parallel and Perpendicular Lines



Figure 2

Since slope measures the steepness of lines, and **parallel lines** have the same steepness, then the **slopes of parallel lines are equal**.

To indicate on a diagram that lines are parallel, we draw on each line arrows pointing in the same direction (see *Figure 2*). To state in mathematical notation that two lines are parallel, we use the \parallel sign.




Figure 3

To see how the slopes of perpendicular lines are related, rotate a line with a given slope $\frac{a}{b}$ (where $b \neq 0$) by 90° , as in *Figure 3*. Observe that under this rotation the vertical change a becomes the horizontal change but in opposite direction ($-a$), and the horizontal change b becomes the vertical change. So, the **slope of the perpendicular line is $-\frac{b}{a}$** . In other words, **slopes of perpendicular lines are opposite reciprocals**. Notice that the **product of perpendicular slopes, $\frac{a}{b} \cdot \left(-\frac{b}{a}\right)$, is equal to -1** .

In the case of $b = 0$, the slope is undefined, so the line is vertical. After rotation by 90° , we obtain a horizontal line, with a slope of zero. So a line with a zero slope and a line with an “undefined” slope can also be considered perpendicular.

To indicate on a diagram that two lines are perpendicular, we draw a square at the intersection of the two lines, as in *Figure 3*. To state in mathematical notation that two lines are perpendicular, we use the \perp sign.

In summary, if m_1 and m_2 are **slopes** of two lines, then the lines are:

- **parallel** iff $m_1 = m_2$, and
- **perpendicular** iff $m_1 = -\frac{1}{m_2}$ (or equivalently, if $m_1 \cdot m_2 = -1$)

In addition, a **horizontal** line (with a slope of **zero**) is **perpendicular** to a **vertical** line (with **undefined** slope).

Example 6 ▶ **Determining Whether the Given Lines are Parallel, Perpendicular, or Neither**

For each pair of linear equations, determine whether the lines are parallel, perpendicular, or neither.

a. $3x + 5y = 7$ $5x - 3y = 4$	b. $y = x$ $2x - 2y = 5$	c. $y = 5$ $y = 5x$
--	------------------------------------	-------------------------------

Solution ▶ **a.** As seen in *Section G1*, the slope of a line given by an equation in standard form, $Ax + By = C$, is equal to $-\frac{A}{B}$. One could confirm this by solving the equation for y and taking the coefficient by x for the slope.

Using this fact, the slope of the line $3x + 5y = 7$ is $-\frac{3}{5}$, and the slope of $5x - 3y = 4$ is $\frac{5}{3}$. Since these two slopes are opposite reciprocals of each other, the two lines are **perpendicular**.


b. The slope of the line $y = x$ is **1** and the slope of $2x - 2y = 5$ is also $\frac{2}{2} = \mathbf{1}$. So, the two lines are parallel.

c. The line $y = 5$ can be seen as $y = 0x + 5$, so its slope is **0**. The slope of the second line, $y = 5x$, is **5**. So, the two lines are neither parallel nor perpendicular.

Collinear Points

Definition 2.2 ▶ Points that lie on the same line are called **collinear**.

Two points are always collinear because there is only one line passing through these points. The question is how could we check if a third point is collinear with the given two points? If we have an equation of the line passing through the first two points, we could plug in the coordinates of the third point and see if the equation is satisfied. If it is, the third point is collinear with the other two. But, can we check if points are collinear without referring to an equation of a line?



Notice that if several points lie on the same line, the slope between any pair of these points will be equal to the slope of this line. So, these slopes will be the same. One can also show that if the slopes between any two points in the group are the same, then such points lie on the same line. So, they are collinear.

Points are **collinear** iff the **slope** between each pair of points is the same.

Example 7 ▶ **Determine Whether the Given Points are Collinear**

Determine whether the points $A(-3,7)$, $B(-1,2)$, and $C = (3,-8)$ are collinear.

Solution ▶ Let m_{AB} represent the slope of \overline{AB} and m_{BC} represent the slope of \overline{BC} . Since

$$m_{AB} = \frac{2-7}{-1-(-3)} = -\frac{5}{2} \quad \text{and} \quad m_{BC} = \frac{-8-2}{3-(-1)} = -\frac{10}{4} = -\frac{5}{2},$$

Then all points A , B , and C lie on the same line. Thus, they are collinear.

Example 8 ▶ **Finding the Missing Coordinate of a Collinear Point**

For what value of y are the points $P(2,2)$, $Q(-1,y)$, and $R(1,6)$ collinear?

Solution ▶ For the points P , Q , and R to be collinear, we need the slopes between any two pairs of these points to be equal. For example, the slope m_{PQ} should be equal to the slope m_{PR} . So, we solve the equation

$$m_{PQ} = m_{PR}$$

for y :

$$\frac{y-2}{-1-2} = \frac{6-2}{1-2}$$

$$\frac{y-2}{-3} = -4$$

$$y-2 = 12$$


$$y = 14$$

Thus, point Q is collinear with points P and R , if $y = 14$.


G.2 Exercises

Given the graph, find the slope of each line.


1.




2.



3.



4.



Given the equation, find the slope of each line.

5. $y = \frac{1}{2}x - 7$

6. $y = -\frac{1}{3}x + 5$

7. $4x - 5y = 2$

8. $3x + 4y = 2$

9. $x = 7$

10. $y = -\frac{3}{4}$

11. $y + x = 1$

12. $-8x - 7y = 24$

13. $-9y - 36 + 4x = 0$

Graph each line satisfying the given information.

14. passing through $(-2, -4)$ with slope $m = 4$

15. passing through $(-1, -2)$ with slope $m = -3$

16. passing through $(-3, 2)$ with slope $m = \frac{1}{2}$

17. passing through $(-3, 4)$ with slope $m = -\frac{2}{5}$

18. passing through $(2, -1)$ with undefined slope

19. passing through $(2, -1)$ with slope $m = 0$

20. Which of the following formulas represent slope?

a. $m = \frac{y_1 - y_2}{x_2 - x_1}$

b. $m = \frac{y_1 - y_2}{x_1 - x_2}$

c. $m = \frac{x_2 - x_1}{y_2 - y_1}$

d. $m = \frac{y_2 - y_1}{x_2 - x_1}$

Find the slope of the line through each pair of points.

21. $(-2, 2), (4, 5)$

22. $(8, 7), (2, -1)$

23. $(9, -4), (3, -8)$

24. $(-5, 2), (-9, 5)$

25. $(-2, 3), (7, -12)$


26. $(3, -1), \left(-\frac{1}{2}, \frac{1}{5}\right)$

27. $(-5, 2), (8, 2)$

28. $(-3, 4), (-3, 10)$





29. $\left(\frac{1}{2}, 6\right), \left(-\frac{2}{3}, \frac{5}{2}\right)$

30. List the line segments in the accompanying figure with respect to their slopes, from the smallest to the largest slope. List the segment with an undefined slope as last.







31. Refer to the graphs of a stock price observed during a 3-month period. Match each situation in **a–d** with the most appropriate graph in **A–D**.

- The price rose sharply during the first month, remained stable during the second month before falling to the original price by the end of the third month.
- The price fell sharply during the first month and then rose slowly for the next two months.
- The price rose sharply during the first month, fell slowly during the second month before rising slowly to its highest peak during the third month.
- The price fell slowly during the first two months before stabilizing in the third month.



Find and interpret the average rate of change illustrated in each graph.



In problems #43-46, sketch a graph depicting each situation. Assume that the roads in each problem are straight.

- 36. The distance that a driver is from home if he starts driving home from a town that is 30 kilometers away and he drives at a constant speed for half an hour.
- 37. The distance that a cyclist is from home if he bikes away from home at 30 kilometers per hour for 30 minutes and then bikes back home at 15 kilometers per hour.
- 38. The distance that Alice is from home if she walks 4 kilometers from home to a shopping centre, stays there for 1.5 hours, and then walks back home. Assume that Alice walks at a constant speed for 30 minutes each way.
- 39. The amount of water in a 500 liters outdoor pool for kids that is filled at the rate of 1500 liters per hour, left full for 4 hours, and then drained at the rate of 3000 liters per hour.

Solve each problem.

- 40. At 6:00 a.m. a 60,000-liter swimming pool was $\frac{1}{3}$ full and at 9:00 a.m. the pool was filled up to $\frac{3}{4}$ of its capacity. Find the rate of filling the pool with the assumption that the rate was constant.
- 41. Jan and Bill plan to drive to Kelowna that is 324 kilometers away. Jan noticed that during one hour they change their location from being $\frac{1}{3}$ of the way to being $\frac{2}{3}$ of the way. Assuming that they drive at a constant rate, what is their average speed of driving?
- 42. Suppose we see a road sign informing that a road grade is 7% for the next 1.5 kilometers. In meters, what would the expected change in elevation be 1.5 kilometers down the road?



Decide whether each pair of lines is parallel, perpendicular, or neither.

- | | | | |
|------------------------------------|---|------------------------------------|--|
| 43. $y = x$
$y = -x$ | 44. $y = 3x - 6$
$y = -\frac{1}{3}x + 5$ | 45. $2x + y = 7$
$-6x - 3y = 1$ | 46. $x = 3$
$x = -2$ |
| 47. $3x + 4y = 3$
$3x - 4y = 5$ | 48. $5x - 2y = 3$
$2x - 5y = 1$ | 49. $y - 4x = 1$
$x + 4y = 3$ | 50. $y = \frac{2}{3}x - 2$
$-2x + 3y = 6$ |

Solve each problem.

- 51. Check whether or not the points $(-2,7)$, $(1,5)$, and $(3,4)$ are collinear.
- 52. The following points, $(2,2)$, $(-1,k)$, and $(1,6)$ are collinear. Find the value of k .

G3

Forms of Linear Equations in Two Variables



Linear equations in two variables can take different forms. Some forms are easier to use for graphing, while others are more suitable for finding an equation of a line given two pieces of information. In this section, we will take a closer look at various forms of linear equations and their utilities.

Forms of Linear Equations

The form of a linear equation that is most useful for graphing lines is the slope-intercept form, as introduced in *Section G1*.

Definition 3.1 ▶ The **slope-intercept form** of the equation of a line with **slope m** and **y -intercept $(0, b)$** is

$$y = mx + b.$$

Example 1 ▶ **Writing and Graphing Equation of a Line in Slope-Intercept Form**

Write the equation in slope-intercept form of the line satisfying the given conditions, and then graph this line.


- slope $-\frac{5}{4}$ and y -intercept $(0, -2)$
- slope $\frac{1}{2}$ and passing through $(-2, -5)$

Solution ▶

- To write this equation, we substitute $m = -\frac{5}{4}$ and $b = -2$ into the slope-intercept form. So, we obtain

$$y = -\frac{5}{4}x - 2.$$

To graph this line, we start with plotting the y -intercept $(0, -2)$. To find the second point, we follow the slope, as in *Example 2, Section G2*. According to the slope $-\frac{5}{4} = \frac{-5}{4}$, starting from $(0, -2)$, we could run 4 units to the right and 5 units down, but then we would go out of the grid. So, this time, let the negative sign in the slope be kept in the denominator, $\frac{5}{-4}$. Thus, we run 4 units to the left and 5 units up to reach the point $(-4, 3)$. Then we draw the line by connecting the two points.



- Since $m = \frac{1}{2}$, our equation has a form $y = \frac{1}{2}x + b$. To find b , we substitute point $(-2, -5)$ into this equation and solve for b . So

$$-5 = \frac{1}{2}(-2) + b$$

gives us


$$-5 = -1 + b$$

and finally

$$b = -4.$$

Therefore, our equation of the line is $y = \frac{1}{2}x - 4$.

We graph it, starting by plotting the given point $(-2, -5)$ and finding the second point by following the slope of $\frac{1}{2}$, as described in *Example 2, Section G2*.



The form of a linear equation that is most useful when writing equations of lines with unknown y -intercept is the slope-point form.

Definition 3.2 ▶ The **slope-point form** of the equation of a line with slope m and passing through the point (x_1, y_1) is

$$y - y_1 = m(x - x_1).$$

This form is based on the defining property of a line. A line can be defined as a set of points with a constant slope m between any two of these points. So, if (x_1, y_1) is a given (fixed) point of the line and (x, y) is any (variable) point of the line, then, since the slope is equal to m for all such points, we can write the equation

$$m = \frac{y - y_1}{x - x_1}.$$

After multiplying by the denominator, we obtain the slope-point formula, as in *Definition 3.2*.

Example 2 ▶ **Writing Equation of a Line Using Slope-Point Form**

Use the slope-point form to write an equation of the line satisfying the given conditions. Leave the answer in the slope-intercept form and then graph the line.

- slope $-\frac{2}{3}$ and passing through $(1, -3)$
- passing through points $(2, 5)$ and $(-1, -2)$

Solution ▶ a. To write this equation, we plug the slope $m = -\frac{2}{3}$ and the coordinates of the point $(1, -3)$ into the slope-point form of a line. So, we obtain


$$y - (-3) = -\frac{2}{3}(x - 1)$$

$$y + 3 = -\frac{2}{3}x + \frac{2}{3}$$

$$y = -\frac{2}{3}x + \frac{2}{3} - \frac{9}{3}$$

$$y = -\frac{2}{3}x - \frac{7}{3}$$

To graph this line, we start with plotting the point $(1, -3)$ and then apply the slope of $-\frac{2}{3}$ to find additional points that belong to the line.



- b. This time the slope is not given, so we will calculate it using the given points, $(2, 5)$ and $(-1, -2)$. Thus,

$$m = \frac{\Delta y}{\Delta x} = \frac{-2 - 5}{-1 - 2} = \frac{-7}{-3} = \frac{7}{3}$$

Then, using the calculated slope and one of the given points, for example $(2, 5)$, we write the slope-point equation of the line


$$y - 5 = \frac{7}{3}(x - 2)$$

and solve it for y :

$$y - 5 = \frac{7}{3}x - \frac{14}{3}$$

$$y = \frac{7}{3}x - \frac{14}{3} + \frac{15}{3}$$

$$y = \frac{7}{3}x + \frac{1}{3}$$



To graph this line, it is enough to connect the two given points.

One of the most popular forms of a linear equation is the standard form. This form is helpful when graphing lines based on x - and y -intercepts, as illustrated in *Example 3, Section G1*.

Definition 3.3 ▶ The **standard form** of a linear equation is

$$Ax + By = C,$$

Where $A, B, C \in \mathbb{R}$, A and B are not both 0, and $A \geq 0$.

When writing linear equations in standard form, the expectation is to use a **nonnegative coefficient A** and **clear any fractions**, if possible. For example, to write $-x + \frac{1}{2}y = 3$ in standard form, we multiply the equation by (-2) , to obtain $2x - y = -6$. In addition, we prefer to write equations in simplest form, where the greatest common factor of A, B , and C is 1. For example, we prefer to write $2x - y = -6$ rather than any multiple of this equation, such as $4x - 2y = -12$, or $6x - 3y = -18$.

Observe that if $B \neq 0$ then the **slope** of the line given by the equation $Ax + By = C$ is $-\frac{A}{B}$.

This is because after solving this equation for y , we obtain $y = -\frac{A}{B}x + \frac{C}{B}$.


If $B = 0$, then the slope is **undefined**, as we are unable to divide by zero.

The form of a linear equation that is most useful when writing equations of lines based on their x - and y -intercepts is the intercept form.

Definition 3.4 ▶ The **intercept form** of a linear equation is

$$\frac{x}{a} + \frac{y}{b} = 1,$$

where $(a, 0)$ is the **x -intercept** and $(0, b)$ is the **y -intercept** of the line.



We should be able to convert a linear equation from one form to another.

Example 3 ▶ **Converting a Linear Equation to a Different Form**

- Write the equation $3x + 7y = 2$ in slope-intercept form.
- Write the equation $y = \frac{3}{5}x + \frac{7}{2}$ in standard form.
- Write the equation $\frac{x}{4} - \frac{y}{3} = 1$ in standard form.

Solution ▶ a. To write the equation $3x + 7y = 2$ in slope-intercept form, we solve it for y .

$$\begin{aligned} 3x + 7y &= 2 \\ 7y &= -3x + 2 \\ y &= -\frac{3}{7}x + \frac{2}{7} \end{aligned}$$

- b. To write the equation $y = \frac{3}{5}x + \frac{7}{2}$ in standard form, we bring the x -term to the left side of the equation and multiply the equation by the LCD, with the appropriate sign.

$$\begin{aligned} y &= \frac{3}{5}x + \frac{7}{2} \\ -\frac{3}{5}x + y &= \frac{7}{2} \\ 6x - 10y &= -35 \end{aligned}$$

- c. To write the equation $\frac{x}{4} - \frac{y}{3} = 1$ in standard form, we multiply it by the LCD, with the appropriate sign.


$$\begin{aligned} \frac{x}{4} - \frac{y}{3} &= 1 \\ 3x - 4y &= 12 \end{aligned}$$

Example 4 ▶ **Writing Equation of a Line Using Intercept Form**

Write an equation of the line passing through points $(0,7)$ and $(-2,0)$. Leave the answer in standard form.

Solution

- ▶ Since point $(0,7)$ is the y -intercept and point $(-2,0)$ is the x -intercept of our line, to write the equation of the line we can use the intercept form with $a = -2$ and $b = 7$. So, we have




$$\frac{x}{-2} + \frac{y}{7} = 1.$$

To change this equation to standard form, we multiply it by the LCD = -14 . Thus,

$$7x - 2y = -14.$$

Equations representing horizontal or vertical lines are special cases of linear equations in standard form, and as such, they deserve special consideration.



The **horizontal line** passing through the point (a, b) has equation $y = b$, while the **vertical line** passing through the same point has equation $x = a$.

The equation of a **horizontal line**, $y = b$, can be shown in standard form as $0x + y = b$. Observe, that the slope of such a line is $-\frac{0}{1} = 0$.

The equation of a **vertical line**, $x = a$, can be shown in standard form as $x + 0y = a$. Observe, that the slope of such a line is $-\frac{1}{0} = \text{undefined}$.


Example 5**▶ Writing Equations of Horizontal and Vertical Lines**

Find equations of the vertical and horizontal lines that pass through the point $(3, -2)$. Then, graph these two lines.

Solution

- ▶ Since x -coordinates of all points of the vertical line, including $(3, -2)$, are the same, then these x -coordinates must be equal to 3. So, the equation of the vertical line is $x = 3$.

Since y -coordinates of all points of a horizontal line, including $(3, -2)$, are the same, then these y -coordinates must be equal to -2 . So, the equation of the horizontal line is $y = -2$.



Here is a summary of the various forms of linear equations.

Forms of Linear Equations		
Equation	Description	When to Use
$y = mx + b$	Slope-Intercept Form slope is m y-intercept is $(0, b)$	This form is ideal for graphing by using the y -intercept and the slope.
$y - y_1 = m(x - x_1)$	Slope-Point Form slope is m the line passes through (x_1, y_1)	This form is ideal for finding the equation of a line if the slope and a point on the line, or two points on the line, are known.

$Ax + By = C$	Standard Form slope is $-\frac{A}{B}$, if $B \neq 0$ x-intercept is $(\frac{C}{A}, 0)$, if $A \neq 0$. y-intercept is $(0, \frac{C}{B})$, if $B \neq 0$.	This form is useful for graphing, as the x- and y-intercepts and the slope can be easily found by dividing appropriate coefficients.
$\frac{x}{a} + \frac{y}{b} = 1$	Intercept Form slope is $-\frac{b}{a}$ x-intercept is $(a, 0)$ y-intercept is $(0, b)$	This form is ideal for graphing, using the x- and y-intercepts.
$y = b$	Horizontal Line slope is 0 y-intercept is $(0, b)$	This form is used to write equations of, for example, horizontal asymptotes.
$x = a$	Vertical Line slope is undefined x-intercept is $(a, 0)$	This form is used to write equations of, for example, vertical asymptotes.

Note: Except for the equations for a horizontal or vertical line, all of the above forms of linear equations can be converted into each other via algebraic transformations.

Writing Equations of Parallel and Perpendicular Lines

Recall that the slopes of parallel lines are the same, and slopes of perpendicular lines are opposite reciprocals. See *Section G2*.

Example 6 ▶ Writing Equations of Parallel Lines Passing Through a Given Point


Find the slope-intercept form of a line parallel to $y = -2x + 5$ that passes through the point $(-4, 5)$. Then, graph both lines on the same grid.

Solution ▶ Since the line is parallel to $y = -2x + 5$, its slope is -2 . So, we plug the slope of -2 and the coordinates of the point $(-4, 5)$ into the slope-point form of a linear equation.

$$y - 5 = -2(x + 4)$$

This can be simplified to the slope-intercept form, as follows:

$$\begin{aligned} y - 5 &= -2x - 8 \\ y &= -2x - 3 \end{aligned}$$



As shown in the accompanying graph, the line $y = -2x - 3$ (in orange) is parallel to the line $y = -2x + 5$ (in green) and passes through the given point $(-4, 5)$.

Example 7 ▶ **Writing Equations of Perpendicular Lines Passing Through a Given Point**

Find the slope-intercept form of a line perpendicular to $2x - 3y = 6$ that passes through the point $(1,4)$. Then, graph both lines on the same grid.


Solution ▶ The slope of the given line, $2x - 3y = 3$, is $\frac{2}{3}$. To find the slope of a perpendicular line, we take the opposite reciprocal of $\frac{2}{3}$, which is $-\frac{3}{2}$. Since we already know the slope and the point, we can plug these pieces of information into the slope-point formula. So, we have

$$y - 4 = -\frac{3}{2}(x - 1)$$

$$y - 4 = -\frac{3}{2}x + \frac{3}{2}$$

$$y = -\frac{3}{2}x + \frac{3}{2} + \frac{8}{2}$$

$$y = -\frac{3}{2}x + \frac{11}{2}$$



As shown in the accompanying graph, the line $2x - 3y = 6$ (in orange) is indeed perpendicular to the line $y = -\frac{3}{2}x + \frac{11}{2}$ (in green) and passes through the given point $(1,4)$.

Linear Equations in Applied Problems

Linear equations can be used to model a variety of applications in sciences, business, and other areas. Here are some examples.

Example 8 ▶ **Given the Rate of Change and the Initial Value, Determine the Linear Model Relating the Variables**


Lucy and Jack bought a sofa for \$1800. They have paid \$300 down and the rest is going to be paid by interest-free monthly payments of \$75 per month, until the bill is paid in full.

- Write an equation to express the amount that is already paid off, A , in terms of the number of months, n , since their purchase.
- Graph the equation found in part a.
- According to the graph, when will the bill be paid in full?

Solution ▶ a. Since each month the couple pays \$75, after n months, the amount paid off by the monthly installments is $75n$. If we add the initial payment of \$300, the equation representing the amount paid off can be written as

$$A = 75n + 300$$

- b. To graph this equation, we use the slope-intercept method. Starting with the A -intercept of 300, we run 1 and rise 75, repeating this process as many times as needed to hit a lattice point on the chosen scale. As indicated in the accompanying graph, some of the points that the line passes through are $(0,300)$, $(4,600)$, $(8,900)$, and $(20,1800)$.



- c. As shown in the graph, \$1800 will be paid off in 20 months.

Example 9

Finding a Linear Equation that Fits the Data Given by Two Ordered Pairs

In Fahrenheit scale, water freezes at 32°F and boils at 212°F . In Celsius scale, water freezes at 0°C and boils at 100°C . Write a linear equation that can be used to calculate the Celsius temperature, C , when the Fahrenheit temperature, F , is known.

Solution



- To predict the Celsius temperature, C , knowing the Fahrenheit temperature, F , we treat the variable C as dependent on the variable F . So, we consider C as the second coordinate when setting up the ordered pairs, (F, C) , of given data. The corresponding freezing temperatures give us the pair $(32,0)$ and the boiling temperatures give us the pair $(212,100)$. To find the equation of a line passing through these two points, first, we calculate the slope, and then, we use the slope-point formula. So, the slope is

$$m = \frac{C_2 - C_1}{F_2 - F_1} = \frac{100 - 0}{212 - 32} = \frac{100}{180} = \frac{5}{9}$$

and using the point $(32,0)$, the equation of the line is

$$C = \frac{5}{9}(F - 32)$$

Example 10

Determining if the Given Set of Data Follows a Linear Pattern

Observe the data given in each table below. Do they follow a linear pattern? If “yes”, find the slope-intercept form of an equation of the line passing through all the given points. If “no”, explain why not.

a.

x	1	3	5	7	9
y	12	15	18	21	24

b.

x	5	10	15	20	25
y	15	21	26	30	35

Solution

- a. The set of points follows a linear pattern if the slopes between consecutive pairs of these points are the same. These slopes are the ratios of increments in y -values to increments in x -values. Notice that the increases between successive x -values of the given points are constantly equal to 2. So, to check if the points follow a linear pattern, it is enough to check if the increases between successive y -values are also constant. Observe that the numbers in the list 12, 15, 18, 21, 24 steadily increase by 3. Thus, the given set of data follow a linear pattern.

To find an equation of the line passing through these points, we use the slope, which is $\frac{3}{2}$, and one of the given points, for example (1,12). By plugging these pieces of information into the slope-point formula, we obtain

$$y - 12 = \frac{3}{2}(x - 1),$$

which after simplifying becomes

$$\begin{aligned} y - 12 &= \frac{3}{2}x - \frac{3}{2} \\ y &= \frac{3}{2}x + \frac{21}{2} \end{aligned}$$

- b. Observe that the increments between consecutive x -values of the given points are constantly equal to 5, while the increments between consecutive y -values in the list 15, 21, 26, 30, 35 are 6, 5, 4, 5. So, they are not constant. Therefore, the given set of data does not follow a linear pattern.

Example 11 ▶ Finding a Linear Model Relating the Number of Items Bought at a Fixed Amount



At a local market, a farmer sells organically grown apples at \$0.50 each and pears at \$0.75 each.

- Write a linear equation in standard form relating the number of apples, a , and pears, p , that can be bought for \$60.
- Graph the equation from part (a).
- Using the graph, find at least 2 points (a, p) satisfying the equation, and interpret their meanings in the context of the problem.

Solution ▶

- a. It costs $0.5a$ dollars to buy a apples. Similarly, it costs $0.75p$ dollars to buy p pears. Since the total charge is \$60, we have

$$0.5a + 0.75p = 60$$


The coefficients can be converted into integers by multiplying the equation by a hundred. This would give us

$$50a + 75p = 6000,$$

which, after dividing by 25, turns into

$$2a + 3p = 240.$$

- b. To graph this equation, we will represent the number of apples, a , on the horizontal axis and the number of pears, p , on the vertical axis, respecting the alphabetical order of labelling the axes. Using the intercept method, we connect points (120,0) and (0,80).



- c. Aside of the intercepts, (120,0) and (0,80), the graph shows us a few more points that satisfy the equation. In particular, (30, 60) and (60, 40) are points of the graph. If a point (a, p) of the graph has integral coefficients, it tells us that \$60 can buy a apples and p pears. For example, the point (30, 60) tells us that **30 apples and 60 pears** can be bought for **\$60**.

G.3 Exercises

Write each equation in **standard form**.

1. $y = -\frac{1}{2}x - 7$

2. $y = \frac{1}{3}x + 5$

3. $\frac{x}{5} + \frac{y}{-4} = 1$

4. $y - 7 = \frac{3}{2}(x - 3)$

5. $y - \frac{5}{2} = -\frac{2}{3}(x + 6)$

6. $2y = -0.21x + 0.35$

Write each equation in **slope-intercept form**.

7. $3y = \frac{1}{2}x - 5$

8. $\frac{x}{3} + \frac{y}{5} = 1$





9. $4x - 5y = 10$

10. $3x + 4y = 7$

11. $y + \frac{3}{2} = \frac{2}{5}(x + 2)$

12. $y - \frac{1}{2} = -\frac{2}{3}\left(x - \frac{1}{2}\right)$

Write an equation in **slope-intercept form** of the line shown in each graph.



Find an equation of the line that satisfies the given conditions. Write the equation in **slope-intercept** and **standard form**.

17. through $(-3, 2)$, with slope $m = \frac{1}{2}$

18. through $(-2, 3)$, with slope $m = -4$

19. with slope $m = \frac{3}{2}$ and y-intercept at -1

20. with slope $m = -\frac{1}{5}$ and y-intercept at 2

21. through $(-1, -2)$, with y-intercept at -3

22. through $(-4, 5)$, with y-intercept at $\frac{3}{2}$

23. through $(2, -1)$ and $(-4, 6)$

24. through $(3, 7)$ and $(-5, 1)$

25. through $\left(-\frac{4}{3}, -2\right)$ and $\left(\frac{4}{5}, \frac{2}{3}\right)$

26. through $\left(\frac{4}{3}, \frac{3}{2}\right)$ and $\left(-\frac{1}{2}, \frac{4}{3}\right)$

Find an equation of the line that satisfies the given conditions.

27. through $(-5, 7)$, with slope 0

28. through $(-2, -4)$, with slope 0

29. through $(-1, -2)$, with undefined slope

30. through $(-3, 4)$, with undefined slope

31. through $(-3, 6)$ and horizontal

32. through $\left(-\frac{5}{3}, -\frac{7}{2}\right)$ and horizontal


33. through $\left(-\frac{3}{4}, -\frac{3}{2}\right)$ and vertical

34. through $(5, -11)$ and vertical


Write an equation in **standard form** for each of the lines described. In each case make a sketch of the given line and the line satisfying the conditions.

35. through (7,2) and parallel to $3x - y = 4$ 36. through (4,1) and parallel to $2x + 5y = 10$
 37. through (-2,3) and parallel to $-x + 2y = 6$ 38. through (-1, -3) and parallel to $-x + 3y = 12$
 39. through (-1,2) and parallel to $y = 3$ 40. through (-1,2) and parallel to $x = -3$
 41. through (6,2) and perpendicular to $2x - y = 5$ 42. through (0,2) and perpendicular to $5x + y = 15$
 43. through (-2,4) and perpendicular to $3x + y = 6$ 44. through (-4, -1) and perpendicular to $x - 3y = 9$
 45. through (3, -4) and perpendicular to $x = 2$ 46. through (3, -4) and perpendicular to $y = -3$

For each situation, write an equation in the form $y = mx + b$, and then answer the question of the problem.

47. Membership in the Apollo Athletic Club costs \$80, plus \$49.95 per month. Express the cost C of the membership in terms of the number of months n that the membership is good for. What is the cost of the one-year membership?
48. A cellphone plan includes 1000 anytime minutes for \$55 per month, plus a one-time activation fee of \$75. Assuming that a cellphone is included in this plan at no additional charge, express the cost C of service in terms of the number of months n of this service. How much would a one-year contract plan cost for this cellphone? 
49. An air compressor can be rented for \$23 per day and a \$60 deposit. Let d represent the number of days that the compressor is rented and C represent the total charge for renting, in dollars.
 a. Write an equation that relates C with d .
 b. Suppose Colin rented the air compressor and paid \$198. For how long did Colin rent the compressor?
50. A car can be rented for \$75 plus \$0.15 per kilometer. Let d represent the number of kilometers driven and C represent the cost of renting, in dollars.
 a. Write an equation that relates C with d .
 b. How many kilometers was the car driven if the total cost of renting is \$101.40?

Solve each problem.

51. Originally there were 8 members of a local high school Math Circle. Three years later, the Math Circle counted 25 members. Assuming that the membership continues to grow at the same rate, find an equation that represents the number N of the Math Circle members t years after.
52. Driving on a highway, Steven noticed a 152-km marker on the side of the road. Ten minutes later, he noticed a 169-km marker. Find a formula that can be used to determine the distance driven d , in kilometers, in terms of the elapsed time t , in hours. 
53. The table below shows the annual tuition and fees at Oxford University for out-of-state students.

Year y	2007	2016
Cost C	\$24400	\$31600

- a. Find the slope-intercept form of a line that fits the given data.
 - b. Interpret the slope in the context of the problem.
 - c. Using the line from (a), find the predicted annual tuition and fees at Oxford University in 2022.
54. The life expectancy for a person born in 1900 was 48 years, and in 2000 it was 77 years. To the nearest year, estimate the life expectancy for someone born in 1970.
55. 3 years after Stan opened his mutual funds account, the amount in the account was \$2540. Two years later, the amount in the account was \$2900. Assuming a constant average increase in \$/year, find a linear equation that represents the amount A in Stan's account t years after it was opened.
56. Connor is a car salesperson in the local auto shop. His pay consists of a base salary and a 1.5% commission on sales. One month, his sales were \$165,000, and his total pay was \$3600.
- a. Write an equation in slope-intercept form that shows Connor's total monthly income I in terms of his monthly sales s .
 - b. Graph the equation developed in (a).
 - c. What does the I -intercept represent in the context of the problem?
 - d. What does the slope represent in the context of the problem?
57. A taxi driver charges \$2.50 as his base fare and a constant amount for each kilometer driven. Helen paid \$7.75 for a 3-kilometer trip.
- a. Find an equation in slope-intercept form that defines the total fare $f(k)$ as a function of the number k of kilometers driven.
 - b. Graph the equation found in (a).
 - c. What does the slope of this graph represent in the above situation?
 - d. How many kilometers were driven if a passenger pays \$23.50?

